

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

2328-124

U.S. Application No. (if known, see 37 CFR 1.5)

09/980585INTERNATIONAL APPLICATION NO.
PCT/PT00/00507INTERNATIONAL FILING DATE
07 June 2000PRIORITY DATE CLAIMED
07 June 1999

TITLE OF INVENTION: A METHOD TO ENABLE ASSESSMENT OF GROWTH AND DEATH OF MICRO-ORGANISMS

APPLICANT(S) FOR DO/EO/US: Esa-Matti LILIUS and Marko VIRTA

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☐ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

ITEMS 11. TO 20. below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☒ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☒ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information: Sequence Listing (10 pages); Five (5) sheets of drawings; Applicant(s) Data Sheet; Statement Pursuant to 37 CFR §1.821(f); and Statement Pursuant to 37 CFR §1.125 with marked-up copy of substitute specification.

U.S. APPLICATION NO. 097/980585 37 CFR 1.50		INTERNATIONAL APPLICATION NO. PCT/F100/00507		ATTORNEY DOCKET NO. 2328-124					
21. <input checked="" type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report Not Prepared by EPO or JPO \$ 1,040.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report has been prepared by the EPO or JPO \$ 890.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$ 740.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but claims did not satisfy provisions of PCT Article 33(1)-(4) \$ 710.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$ 100.00				<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 80%;">CALCULATIONS</td> <td style="width: 20%; text-align: center;">PTO USE ONLY</td> </tr> <tr> <td colspan="2" style="height: 150px; vertical-align: top;"> <div style="text-align: right; margin-top: 10px;"> ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 1,040.00 </div> </td> </tr> </table>		CALCULATIONS	PTO USE ONLY	<div style="text-align: right; margin-top: 10px;"> ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 1,040.00 </div>	
CALCULATIONS	PTO USE ONLY								
<div style="text-align: right; margin-top: 10px;"> ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 1,040.00 </div>									
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$					
Claims	Number Filed	Number Extra	Rate						
Total Claims	10 -20 =	0	X \$18.00	\$00.00					
Independent Claims	01 -3 =	0	X \$84.00	\$00.00					
Multiple dependent claim(s) (if applicable)				+\$280.00	\$00.00				
TOTAL OF ABOVE CALCULATIONS =				\$1,040.00					
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$520.00					
SUBTOTAL =				\$520.00					
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$					
TOTAL NATIONAL FEE =				\$520.00					
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				\$					
TOTAL FEES ENCLOSED =				\$520.00					
				Amount to be refunded	\$				
				charged	\$				
a. <input checked="" type="checkbox"/> A check in the amount of \$520.00 to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. 02-2135 in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-2135. A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.									
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.									
SEND ALL CORRESPONDENCE TO:									
Signature:									
Customer No. 6449									
Name: <u>Jeffrey L. Ihnen</u>									
Registration Number: <u>28.957</u>									
Jeffrey L. Ihnen Rodwell, Figg, Ernst & Manbeck 555 13th St., N.W. Washington, D.C. 20004 Phone: 202/783-6040									


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE	<i>Application Number</i>	New Application
	<i>Filing Date</i>	Herewith
	<i>First Named Inventor</i>	Esa-Matti LILJUS
	<i>Group Art Unit</i>	Unassigned
	<i>Examiner Name</i>	Unassigned
	<i>Attorney Docket No.</i>	2328-124
<i>Title of the Invention:</i> A METHOD TO ENABLE ASSESSMENT OF GROWTH AND DEATH OF MICRO-ORGANISMS		

STATEMENT PURSUANT TO 37 CFR 1.125(b)

Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

In the matter of the above-identified nonprovisional application, filed concurrently herewith, Applicants are submitting a substitute copy of the specification. The substitute specification contains no new matter. A marked-up version of the substitute specification, in which additions are indicated by underlining and deletions are indicated by brackets, is attached.

RESPECTFULLY SUBMITTED,					
NAME AND REG. NUMBER	Jeffrey L. Ihnen, Reg. No. 28,957				
SIGNATURE		DATE	04 December 2001		
Address	ROTHWELL, FIGG, ERNST & MANBECK, pc Suite 701-East, 555 13th Street, N.W.				
City	Washington	State	D.C.	Zip Code	20004
Country	U.S.A.	Telephone	202-783-6040	Fax	202-783-6031

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE	<i>Application Number</i>	09/980,585
	<i>Filing Date</i>	4 December 2001
	<i>First Named Inventor</i>	Esa-Matti LILJUS
	<i>Group Art Unit</i>	Unassigned
	<i>Examiner Name</i>	Unassigned
	<i>Attorney Docket No.</i>	2328-124
<i>Title of the Invention:</i> A METHOD TO ENABLE ASSESSMENT OF GROWTH AND DEATH OF MICRO-ORGANISMS		

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

In response to the Notice of Missing Requirements, response copy attached, please amend the above-identified application as follows.

IN THE SEQUENCE LISTING:

Please replace the originally filed Sequence Listing with the substitute Sequence Listing attached hereto.

REMARKS


The Sequence Listing has been revised in view of the inability of the programs to correctly number and accept a nucleic acid sequence which contains more than one coding sequence. Consequently, each sequence originally identified as "CDS" in SEQ ID NO:1 of the original Sequence Listing have been relabeled as "gene" to prevent any errors with the Checker software. The proteins encoded by these "genes" are set forth in SEQ ID Nos:2-4 as in the original Sequence Listing. The substitute Sequence Listing contains no new matter and its entry is requested.

Serial No.: 09/980,585

04 March 2002

Page 2

A computer readable copy of the Sequence Listing is submitted concurrently herewith.

RESPECTFULLY SUBMITTED,			
NAME AND REG. NUMBER	Jeffrey L. Ihnen, Reg. No. 28,957		
SIGNATURE		DATE	04 March 2002

Address	ROTHWELL, FIGG, ERNST & MANBECK, pc 1425 K Street, N.W., Suite 800				
City	Washington	State	D.C.	Zip Code	20005
Country	U.S.A.	Telephone	202-783-6040	Fax	202-783-6031

Attachment: substitute Sequence Listing

09/980,585

TITLE OF THE INVENTION

A METHOD TO ENABLE ASSESSMENT OF
GROWTH AND DEATH OF MICRO-ORGANISMS

5 CROSS-REFERENCE TO RELATED APPLICATION

[0001] The present application is a national stage filing under 35 U.S.C. §371 of PCT/FI00/00507 filed on 7 June 2000 and claims priority under 35 U.S.C. §119 to Finland patent application No. 991296 filed on 7 June 1999.

10 [0002] This invention relates to a method to enable the assessment of growth and death of a micro-organism within a chosen time period in an environment of interest.

BACKGROUND OF THE INVENTION

[0003] When studying growth and death of a micro-organism under the influence of specific environments, e.g. production and storage environments that e.g. could or could not be refrigerated, or involving chemicals or matrixes, e.g. antibiotics, microbial toxins, heavy metals and serum complement, microbial cultures are most often incubated for hours or days. In these circumstances death and growth occur simultaneously. If additionally some of the cells lyse, e.g. when analysing the serum complement, it is difficult to know to what one should compare the amount of living cells at the end of the experiment. Convenient methods to determine the number of living cells, e.g. by measuring luciferase bioluminescence, are known but if no more information is available it is impossible to assess to what extent growth or/and death of the micro-organisms takes or has taken place.

25 [0004] Growth rates and death rates of micro-organisms in specific environments are of interest in many areas. Death rates and growth rates of micro-organisms and especially harmful and/or pathogenic micro-organisms are of importance in risk assessments of products of the pharmaceutical industry and products for human consumption with regard to their production, storage and distribution to the consumers. Knowledge of death and growth rates of micro-organisms are of particular importance in specific applications such as in the development of
30 antibiotics, disinfectants and bactericidal products or monitoring of sterilisation, disinfection and cleaning processes.

[0005] Reporter genes coding for luminescent or/and fluorescent products have been introduced to micro-organisms to enable the assessment of the quantity or survival of living

micro-organisms (WO 96/23898, WO 98/14605, WO 98/30715, WO 98/36081, US 5,824,468). Even simultaneous use of luminescent and fluorescent markers has been used (Fratamico et al., Journal of Food Protection, Vol 50 No 10, 1997, 1167-1173). Luminescent and fluorescent markers have, however, only been used as markers for survival of micro-organisms and the use of two different markers within one micro-organism enabling the differentiation between growth and death rates has not been reported.

OBJECT AND SUMMARY OF THE INVENTION

[0006] The object of the present invention is to provide a method to enable the assessment of the growth and death of a micro-organism within a chosen time period in an environment of interest by introducing into said micro-organism at least two reporter genes. The method is characterised in that

- a) said reporter genes code for luminescent and/or fluorescent products and within said time period and environment at least two said products of the following are produced:
 - i) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that are or have been alive within said chosen time period,
 - ii) a product present in said environment of interest in an essentially known proportion to the amount of cells alive at any moment within said chosen time period and
 - iii) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that have died within said chosen time period,
 and said products can be measured through their luminescence and/or fluorescence;
- b) said micro-organism is incubated within the environment of interest and said luminescence and/or fluorescence is detected after said chosen time period, and
- c) the growth and death rate of the said micro-organism is assessed based on at least two of the following:

- i) the known proportion of luminescence or fluorescence to the amount of cells alive after any said chosen time period,
- ii) the known proportion of luminescence or fluorescence to the total amount of cells that are or have been alive within any said chosen time period, and
- iii) the known proportion of luminescence or fluorescence to the total amount of cells that have died within any said chosen time period.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007] Figure 1 shows plasmid pGFP+luc* including genes for both GFP and firefly luciferase.

[0008] Figure 2 shows fluorescence during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0009] Figure 3 shows luminescence during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0010] Figure 4 shows the amount of living cells, i.e. colony forming units, according to plating during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0011] Figure 5 shows the percentage of living cells according to live/dead staining and flow cytometric analysis during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0012] Figure 6 shows fluorescence before and after incubation with serum complement during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of serum complement in the cell culture.

[0013] Figure 7 shows luminescence before and after incubation with serum complement during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of serum complement in the cell culture.

[0014] Figure 8 shows the percentage of living cells according to plating during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of serum complement in the cell culture.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The method according to the present invention can be used to assess the growth and death rate of a micro-organism within a chosen time period in any particular environment of interest. The method is applicable if two different marker genes can be introduced to the micro-organism that code for luminescent and/or fluorescent products, and the products of these fulfil at least two of the following three criteria:

a) a said luminescent product luminesces or said fluorescent product fluoresces in an essentially known proportion to the amount of cells of said micro-organism alive within said chosen time period ;

b) a said luminescent product luminesces or said fluorescent product fluoresces in an essentially known proportion to the amount of cells of said micro-organism that are or have been alive within said chosen time period, and

c) a said luminescent product luminesces or said fluorescent product fluoresces in an essentially known proportion to the amount of cells of said micro-organism that have died within said chosen time period.

[0016] In the present application the concept "micro-organism" means any micro-organism into which marker genes can be introduced so, that they will function according to the invention. "Micro-organism" can therefore stand for bacteria, yeast or fungi.

[0017] The concept of "introducing a marker gene into a micro-organism" means any method by which a marker gene can be made to function within the micro-organism according to the invention. One way of introducing marker genes into micro-organism is by constructing a recombinant strain. This can be done by transforming a strain with a plasmid including the marker genes. An alternative way to introduce reporter genes to bacteria is to utilise transposable elements. In this technique, reporter genes are inserted between insertion sequences in a delivery plasmid. The plasmid is then introduced to a cell by e.g. conjugation of transformation, and once inside the cell, genes surrounded by the insertion sequences are integrated into bacterial chromosome. Integration is stable, i.e. there is no need for a selectable marker such as antibiotic resistance.

[0018] Assessment of the growth rate and death rate of a micro-organism can be of interest in many specific environments. Within pharmaceutical research the effect of different drugs and candidates for drugs, e.g. antibiotics, on the survival of pathogenic, but also the

beneficial micro-organisms of the gut, is of interest. Thus the ultimate interest is in the behaviour of these micro-organisms in a physiological environment affected by drugs.

[0019] Another vast area where the possibility of assessing growth and death rate of specific micro-organisms is of interest is that of production, processing, storage and distribution of all products for human consumption. In this area the behaviour of pathogenic or potentially harmful micro-organisms in the different environments of the life cycle of these products is of special interest and involves many different aspects such as the influence of temperature, humidity or light and the possible use of preservatives etc.

[0020] Additionally growth and death rates of micro-organisms can be of interest for environmental evaluations e.g. when evaluating the effect of emissions into the environment.

[0021] Luminescent or fluorescent products coded by reporter genes in different embodiments of this invention can vary as long as their proportion to either the total amount of cells alive, to cells that are or have been alive, or to cells that have died is essentially known. Growth and death rate can be assessed if two of the following: cells alive, cells that are or have been alive, or cells that have died can be determined. Thus luminescence and/or fluorescence measured can be e.g. of a product which is expressed e.g. constitutively or triggered by a specific phase (e.g. replication or death) of the lifecycle of each cell, is stable or labile or which luminescence or fluorescence is dependant on a factor that relates e.g. to a specific phase of the lifecycle of each cell. Depending on the individual characteristics of said product—how produced, stable or labile, possible dependence of its luminescence or fluorescence of said factors etc.—the measured luminescence or fluorescence can be in proportion to one of the three unknown of which two must be known to be able to assess the growth rate and death rate of said cells.

[0022] According to one specific embodiment of the invention assessment of the growth and death rate of an *Esherichia coli* strain under the influence of different chemicals or matrixes was enabled by constructing a recombinant strain, which expresses both luciferase and GFP. Altogether the effect of a number of different chemicals and matrixes, such as CdCl_2 , ethanol, the antibiotics chloramphenicol, rifampicin, and tetracyclin, as well as serum complement on said recombinant *E. coli* strain was tested and found applicable.

[0023] The invention will be described in more detail by the following study in which the growth rate and death rate of a recombinant *Esherichia coli* strain, which expresses both luciferase and GFP, is assessed under the influence of ethanol or serum complement.

Summary of the study

[0024] Genes for luciferase and green fluorescent protein have recently raised interest as reporter genes. Luciferase is an enzyme that produces luminescence in the presence of substrate luciferin, molecular oxygen and ATP. Green fluorescent protein (GFP), produces green fluorescence when excited with light. Many mutated forms of GFP have been introduced: some have different excitation and emission wavelengths from the wild type and some mutants form more stable proteins at higher temperatures.

[0025] We constructed a recombinant strain of *E. coli*, which expresses both luciferase and GFP. In our construction we used a mutant of GFP, which is more stable at temperatures over +30 °C and it matures quicker than the wild type. Luciferase was from North American firefly, *Photinus pyralis*.

[0026] The *E. coli* strain MC1061 was transformed with a plasmid including genes for both GFP and firefly luciferase. Figure 1 describes the plasmid in general. The sequence of the plasmid is disclosed in the sequence listing. Essential codings of the sequence are as follows:

lac promoter	95–199
GFP	289–1008
firefly luciferase	1044–2696
β -lactamase	3251–4111

[0027] In our construct, see Figure 1, the luciferase gene is situated next to the GFP gene and both genes are transcribed in the same direction. The transcription is started at the lac promoter in front of GFP. The lac promoter is constitutively active, because the MC1061 cells lack its repressor. The plasmid also has a gene for ampicillin resistance (β -lactamase).

[0028] The transformed *E. coli* strain was propagated under the influence of different concentrations of ethanol or serum complement.

Methods

Growth conditions

[0029] One colony from a pure culture plate was inoculated in 5 ml of LB-medium with ampicillin (100 μ g/ml) and grown at +37 °C in a shaker, 250 rpm, for about three hours. After that, the number of cells per millilitre was determined with flow cytometry by using fluorescent spherical latex particles as a reference. One million cells were then removed to an erlenmeyer

with 50 ml of LB medium and ampicillin. The culture was grown over night in a shaker, 190 rpm, at room temperature to prevent the culture from growing into the stationary phase during the night. In the morning, the culture was transferred to and grown in a shaker, 330 rpm, until the stationary phase was reached or used after growth at +30 °C for about 1 h to study the influence of ethanol or serum complement as described below.

Influence of chemicals on the propagation of *E. coli*

[0030] The culture obtained as described above was used to study the influence of ethanol or serum complement as follows:

Ethanol

[0031] Ethanol (Aa, Primalco Oy) was diluted into pure water to obtain concentrations of 50, 45, 25, 10, 5, 1 and 0 % of ethanol when 500 µl of said dilution was added to 500 µl of said culture in an eppendorf tube. The mixture was vortexed and incubated for 5 minutes before measuring fluorescence and luminescence. Live cells were again counted by plating and also by live/dead staining. In the live/dead protocol used the stain *cyto 9* stains all cells whereas propidium iodide stains only the dead cells. After staining, cells are passed through a flow cytometer, with which dead and live cells can be differentiated and their proportion determined. (Virta et al. (1998) Appl. Environ. Microbiol. 64: 515–519.)

Serum complement

[0032] The influence of serum complement on the said recombinant *E. coli* strain was studied using an incubation time of 90 min as described for a different recombinant *E. coli* strain used in Virta et al. (1998) Appl. Environ. Microbiol. 64: 515–519.

Fluorescence and luminescence measurements

[0033] The measurements were done with a combined fluoro- and luminometer, Fluoroscan Ascent FL, provided by Labsystems Ltd. (Helsinki, Finland). Cell growth was simultaneously followed with a flow cytometer.

[0034] For the measurements, 100 µl of bacterial culture was pipetted into the microtiter plate wells. Fluorescence was measured using 485 nm for excitation and 510 nm for emission. Measuring time was 20 ms. After the fluorescence measurement 100 µl of luciferin in 0.1 M citric acid-sodium citrate buffer (pH 5.0) was dispensed into the wells and the plate was shaken

for two minutes (shaking diameter 1 mm, 1 020 rpm), after which luminescence was recorded with a measuring time of 1000 ms.

Plating

[0035] Samples for plating were diluted 10^2 to 10^7 fold with 150 mM NaCl and plated onto L agar plates (L broth containing 1.6 % agar). Colonies were counted after overnight incubation at 37 °C.

Live/dead staining and Flow cytometric analysis

[0036] Bacteria from 1 000 µl of cell culture were used for live/dead staining and flow cytometric analysis using a LIVE/DEAD BacLight bacterial viability kit (catalogue no. L-7005) for microscopy and quantitative analysis obtained from Molecular Probes Europe (Leiden, The Netherlands) and Fluoresbrite beads (diameter, 1.8 µm) obtained from Polysciences Inc. (Warrington, Pa.) as described in Virta et al. (1998) Appl. Environ. Microbiol. 64: 515–519.

Results

[0037] When the cultures were transferred to +30 °C, the cells grew logarithmically for 1-4 hours depending on the initial cell concentration. Luminescence and fluorescence rose logarithmically and were essentially constant per cell. Thus cell number could be assessed based on luminescence or fluorescence.

[0038] When ethanol was added in different concentrations to the growth medium (see Figures 4 and 5) death was, after a very short incubation period of 5 min, more or less insignificant at ethanol concentrations below 5 % and became more significant with increasing ethanol concentration reaching very pronounced significance at ethanol concentrations above 10 %. Correspondingly fluorescence (Figure 2) was essentially constant whatever the ethanol concentration in spite of dramatically decreasing corresponding live cell count according to plate count (Figure 4) and percentage of live cells according to the live/dead staining (Figure 5) whereas luminescence (Figure 3) dropped dramatically essentially corresponding to the dramatic drop in plate count (Figure 4) and the percentage of live cells (Figure 5) with increased ethanol concentration.

[0039] The effect of serum complement on the growth and death of *E. coli* is shown in Figures 6 to 8. Fluorescence (Figure 6) and luminescence (Figure 7) are shown before (squares) and after (circles) incubation for 90 minutes with serum complement. Fluorescence (Figure 6) is

slightly increased, during incubation regardless of the concentration of serum, whereas luminescence (Figure 7) decreases during incubation with increasing serum concentration. The decrease of luminescence during incubation with increasing concentrations of serum correlates clearly with the percentage of cells alive after incubation (Figure 8).

CLAIMS

1. A method to enable the assessment of the growth rate and death rate of a micro-organism within a chosen time period in an environment of interest by introducing into said micro-organism at least two reporter genes, which method is characterised in that
 - a) said reporter genes code for luminescent and/or fluorescent products and within said time period and environment at least two said products of the following are produced:
 - i) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that are or have been alive within said chosen time period,
 - ii) a product present in said environment of interest in an essentially known proportion to the amount of cells alive at any moment within said chosen time period, and
 - iii) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that have died within said chosen time period,and said products can be measured through their luminescence and/or fluorescence;
 - b) said micro-organism is incubated within the environment of interest and said luminescence and/or fluorescence is detected after said chosen time period, and
 - c) the growth and death rate of the said micro-organism is assessed based on at least two of the following:
 - i) the known proportion of luminescence or fluorescence to the amount of cells alive after any said chosen time period,
 - ii) the known proportion of luminescence or fluorescence to the total amount of cells that are or have been alive within any said chosen time period, and
 - iii) the known proportion of luminescence or fluorescence to the total amount of cells that have died within any said chosen time period.
2. The method according to claim 1 characterised in that said micro-organism is a gram negative bacteria, e.g. *Escherichia coli*.

3. The method according to claim 1 [or 2] characterised in that
- a) one reporter gene coding for a luminescent product is luciferase, which is used for the determination of amount of cells alive at any moment within said chosen time period, and
 - b) another reporter gene coding for a fluorescent product is green fluorescent protein (GFP), which is used for the determination of total amount of cells of said micro-organism that are or have been alive within said chosen time period.
4. The method according to claim 1 [or 2] characterised in that said reporter genes are introduced into said micro-organism in a plasmid.
5. The [A] method according to [the methods of] claim 3 [or 4] characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).
6. The method according to claim 2 characterised in that
- a) one reporter gene coding for a luminescent product is luciferase, which is used for the determination of amount of cells alive at any moment within said chosen time period, and
 - b) another reporter gene coding for a fluorescent product is green fluorescent protein (GFP), which is used for the determination of total amount of cells of said micro-organism that are or have been alive within said chosen time period.
7. The method according to claim 2 characterised in that said reporter genes are introduced into said micro-organism in a plasmid.
8. The method according to claim 4 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).
9. The method according to claim 6 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).

10. The method according to claim 7 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).

204020.5850660

ABSTRACT

[0040] A method to enable the assessment of the growth rate and death rate of a micro-organism within a chosen time period in an environment of interest. The method is characterised in that

a) two reporter genes are introduced to said micro-organism wherein, the reporter genes used code for luminescent and/or fluorescent products and at least two of the following products: an essentially stable product produced in an essentially known proportion to the total amount of cells of said micro-organism that are or have been alive within said chosen time period; a product present in an essentially known proportion to the amount of cells alive at any moment within said chosen time period; and an essentially stable product produced in an essentially known proportion to the total amount of cells of the said micro-organism that have died within said chosen time period, and said products can be measured through their luminescence and/or fluorescence;

b) the said micro-organism is incubated and said luminescence and/or fluorescence is detected after said chosen time periods, and

c) the growth and death rate of the said micro-organism is assessed based on at least two of the following: the known proportion of luminescence or fluorescence to the amount of cells alive after any said chosen time period; the known proportion of luminescence or fluorescence to the total amount of cells that are or have been alive within any said chosen time period; and the known proportion of luminescence or fluorescence to the total amount of cells that have died within any said chosen time period.

5/pv1b

TITLE OF THE INVENTION

A METHOD TO ENABLE ASSESSMENT OF GROWTH AND DEATH OF MICRO-ORGANISMS

5 CROSS-REFERENCE TO RELATED APPLICATION

[0001] The present application is a national stage filing under 35 U.S.C. §371 of PCT/FI00/00507 filed on 7 June 2000 and claims priority under 35 U.S.C. §119 to Finland patent application No. 991296 filed on 7 June 1999.

10 [0002] This invention relates to a method to enable the assessment of growth and death of a micro-organism within a chosen time period in an environment of interest.

BACKGROUND OF THE INVENTION

[0003] When studying growth and death of a micro-organism under the influence of specific environments, e.g. production and storage environments that e.g. could or could not be refrigerated, or involving chemicals or matrixes, e.g. antibiotics, microbial toxins, heavy metals and serum complement, microbial cultures are most often incubated for hours or days. In these circumstances death and growth occur simultaneously. If additionally some of the cells lyse, e.g. when analysing the serum complement, it is difficult to know to what one should compare the amount of living cells at the end of the experiment. Convenient methods to determine the number of living cells, e.g. by measuring luciferase bioluminescence, are known but if no more information is available it is impossible to assess to what extent growth or/and death of the micro-organisms takes or has taken place.

25 [0004] Growth rates and death rates of micro-organisms in specific environments are of interest in many areas. Death rates and growth rates of micro-organisms and especially harmful and/or pathogenic micro-organisms are of importance in risk assessments of products of the pharmaceutical industry and products for human consumption with regard to their production, storage and distribution to the consumers. Knowledge of death and growth rates of micro-organisms are of particular importance in specific applications such as in the development of
30 antibiotics, disinfectants and bactericidal products or monitoring of sterilisation, disinfection and cleaning processes.

[0005] Reporter genes coding for luminescent or/and fluorescent products have been introduced to micro-organisms to enable the assessment of the quantity or survival of living

0365555-03072

micro-organisms (WO 96/23898, WO 98/14605, WO 98/30715, WO 98/36081, US 5,824,468). Even simultaneous use of luminescent and fluorescent markers has been used (Fratamico et al., Journal of Food Protection, Vol 50 No 10, 1997, 1167-1173). Luminescent and fluorescent markers have, however, only been used as markers for survival of micro-organisms and the use of two different markers within one micro-organism enabling the differentiation between growth and death rates has not been reported.

OBJECT AND SUMMARY OF THE INVENTION

[0006] The object of the present invention is to provide a method to enable the assessment of the growth and death of a micro-organism within a chosen time period in an environment of interest by introducing into said micro-organism at least two reporter genes. The method is characterised in that

- a) said reporter genes code for luminescent and/or fluorescent products and within said time period and environment at least two said products of the following are produced:
 - i) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that are or have been alive within said chosen time period,
 - ii) a product present in said environment of interest in an essentially known proportion to the amount of cells alive at any moment within said chosen time period and
 - iii) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that have died within said chosen time period,
 and said products can be measured through their luminescence and/or fluorescence;
- b) said micro-organism is incubated within the environment of interest and said luminescence and/or fluorescence is detected after said chosen time period, and
- c) the growth and death rate of the said micro-organism is assessed based on at least two of the following:

- i) the known proportion of luminescence or fluorescence to the amount of cells alive after any said chosen time period,
- ii) the known proportion of luminescence or fluorescence to the total amount of cells that are or have been alive within any said chosen time period, and
- iii) the known proportion of luminescence or fluorescence to the total amount of cells that have died within any said chosen time period.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007] Figure 1 shows plasmid pGFP+luc* including genes for both GFP and firefly luciferase.

[0008] Figure 2 shows fluorescence during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0009] Figure 3 shows luminescence during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0010] Figure 4 shows the amount of living cells, i.e. colony forming units, according to plating during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0011] Figure 5 shows the percentage of living cells according to live/dead staining and flow cytometric analysis during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of ethanol in the cell culture.

[0012] Figure 6 shows fluorescence before and after incubation with serum complement during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of serum complement in the cell culture.

[0013] Figure 7 shows luminescence before and after incubation with serum complement during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of serum complement in the cell culture.

[0014] Figure 8 shows the percentage of living cells according to plating during growth phase of *E. coli* with plasmid pGFP+luc* at 30 °C as a function of the concentration of serum complement in the cell culture.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The method according to the present invention can be used to assess the growth and death rate of a micro-organism within a chosen time period in any particular environment of interest. The method is applicable if two different marker genes can be introduced to the micro-organism that code for luminescent and/or fluorescent products, and the products of these fulfil at least two of the following three criteria:

a) a said luminescent product luminesces or said fluorescent product fluoresces in an essentially known proportion to the amount of cells of said micro-organism alive within said chosen time period ;

b) a said luminescent product luminesces or said fluorescent product fluoresces in an essentially known proportion to the amount of cells of said micro-organism that are or have been alive within said chosen time period, and

c) a said luminescent product luminesces or said fluorescent product fluoresces in an essentially known proportion to the amount of cells of said micro-organism that have died within said chosen time period.

[0016] In the present application the concept "micro-organism" means any micro-organism into which marker genes can be introduced so, that they will function according to the invention. "Micro-organism" can therefore stand for bacteria, yeast or fungi.

[0017] The concept of "introducing a marker gene into a micro-organism" means any method by which a marker gene can be made to function within the micro-organism according to the invention. One way of introducing marker genes into micro-organism is by constructing a recombinant strain. This can be done by transforming a strain with a plasmid including the marker genes. An alternative way to introduce reporter genes to bacteria is to utilise transposable elements. In this technique, reporter genes are inserted between insertion sequences in a delivery plasmid. The plasmid is then introduced to a cell by e.g. conjugation of transformation, and once inside the cell, genes surrounded by the insertion sequences are integrated into bacterial chromosome. Integration is stable, i.e. there is no need for a selectable marker such as antibiotic resistance.

[0018] Assessment of the growth rate and death rate of a micro-organism can be of interest in many specific environments. Within pharmaceutical research the effect of different drugs and candidates for drugs, e.g. antibiotics, on the survival of pathogenic, but also the

beneficial micro-organisms of the gut, is of interest. Thus the ultimate interest is in the behaviour of these micro-organisms in a physiological environment affected by drugs.

[0019] Another vast area where the possibility of assessing growth and death rate of specific micro-organisms is of interest is that of production, processing, storage and distribution of all products for human consumption. In this area the behaviour of pathogenic or potentially harmful micro-organisms in the different environments of the life cycle of these products is of special interest and involves many different aspects such as the influence of temperature, humidity or light and the possible use of preservatives etc.

[0020] Additionally growth and death rates of micro-organisms can be of interest for environmental evaluations e.g. when evaluating the effect of emissions into the environment.

[0021] Luminescent or fluorescent products coded by reporter genes in different embodiments of this invention can vary as long as their proportion to either the total amount of cells alive, to cells that are or have been alive, or to cells that have died is essentially known. Growth and death rate can be assessed if two of the following: cells alive, cells that are or have been alive, or cells that have died can be determined. Thus luminescence and/or fluorescence measured can be e.g. of a product which is expressed e.g. constitutively or triggered by a specific phase (e.g. replication or death) of the lifecycle of each cell, is stable or labile or which luminescence or fluorescence is dependant on a factor that relates e.g. to a specific phase of the lifecycle of each cell. Depending on the individual characteristics of said product—how produced, stable or labile, possible dependence of its luminescence or fluorescence of said factors etc.—the measured luminescence or fluorescence can be in proportion to one of the three unknown of which two must be known to be able to assess the growth rate and death rate of said cells.

[0022] According to one specific embodiment of the invention assessment of the growth and death rate of an *Escherichia coli* strain under the influence of different chemicals or matrixes was enabled by constructing a recombinant strain, which expresses both luciferase and GFP. Altogether the effect of a number of different chemicals and matrixes, such as CdCl₂, ethanol, the antibiotics chloramphenicol, rifampicin, and tetracyclin, as well as serum complement on said recombinant *E. coli* strain was tested and found applicable.

[0023] The invention will be described in more detail by the following study in which the growth rate and death rate of a recombinant *Escherichia coli* strain, which expresses both luciferase and GFP, is assessed under the influence of ethanol or serum complement.

Summary of the study

[0024] Genes for luciferase and green fluorescent protein have recently raised interest as reporter genes. Luciferase is an enzyme that produces luminescence in the presence of substrate luciferin, molecular oxygen and ATP. Green fluorescent protein (GFP), produces green fluorescence when excited with light. Many mutated forms of GFP have been introduced: some have different excitation and emission wavelengths from the wild type and some mutants form more stable proteins at higher temperatures.

[0025] We constructed a recombinant strain of *E. coli*, which expresses both luciferase and GFP. In our construction we used a mutant of GFP, which is more stable at temperatures over +30 °C and it matures quicker than the wild type. Luciferase was from North American firefly, *Photinus pyralis*.

[0026] The *E. coli* strain MC1061 was transformed with a plasmid including genes for both GFP and firefly luciferase. Figure 1 describes the plasmid in general. The sequence of the plasmid is disclosed in the sequence listing. Essential codings of the sequence are as follows:

lac promoter	95–199
GFP	289–1008
firefly luciferase	1044–2696
β-lactamase	3251–4111

[0027] In our construct, see Figure 1, the luciferase gene is situated next to the GFP gene and both genes are transcribed in the same direction. The transcription is started at the lac promoter in front of GFP. The lac promoter is constitutively active, because the MC1061 cells lack its repressor. The plasmid also has a gene for ampicillin resistance (β-lactamase).

[0028] The transformed *E. coli* strain was propagated under the influence of different concentrations of ethanol or serum complement.

Methods

Growth conditions

[0029] One colony from a pure culture plate was inoculated in 5 ml of LB-medium with ampicillin (100 µg/ml) and grown at +37 °C in a shaker, 250 rpm, for about three hours. After that, the number of cells per millilitre was determined with flow cytometry by using fluorescent spherical latex particles as a reference. One million cells were then removed to an erlenmeyer

with 50 ml of LB medium and ampicillin. The culture was grown over night in a shaker, 190 rpm, at room temperature to prevent the culture from growing into the stationary phase during the night. In the morning, the culture was transferred to and grown in a shaker, 330 rpm, until the stationary phase was reached or used after growth at +30 °C for about 1 h to study the influence of ethanol or serum complement as described below.

Influence of chemicals on the propagation of *E. coli*

[0030] The culture obtained as described above was used to study the influence of ethanol or serum complement as follows:

Ethanol

[0031] Ethanol (Aa, Primalco Oy) was diluted into pure water to obtain concentrations of 50, 45, 25, 10, 5, 1 and 0 % of ethanol when 500 µl of said dilution was added to 500 µl of said culture in an eppendorf tube. The mixture was vortexed and incubated for 5 minutes before measuring fluorescence and luminescence. Live cells were again counted by plating and also by live/dead staining. In the live/dead protocol used the stain *cyto 9* stains all cells whereas propidium iodide stains only the dead cells. After staining, cells are passed through a flow cytometer, with which dead and live cells can be differentiated and their proportion determined. (Virta et al. (1998) Appl. Environ. Microbiol. 64: 515–519.)

Serum complement

[0032] The influence of serum complement on the said recombinant *E. coli* strain was studied using an incubation time of 90 min as described for a different recombinant *E. coli* strain used in Virta et al. (1998) Appl. Environ. Microbiol. 64: 515–519.

Fluorescence and luminescence measurements

[0033] The measurements were done with a combined fluoro- and luminometer, Fluoroscanner Ascent FL, provided by Labsystems Ltd. (Helsinki, Finland). Cell growth was simultaneously followed with a flow cytometer.

[0034] For the measurements, 100 µl of bacterial culture was pipetted into the microtiter plate wells. Fluorescence was measured using 485 nm for excitation and 510 nm for emission. Measuring time was 20 ms. After the fluorescence measurement 100 µl of luciferin in 0.1 M citric acid-sodium citrate buffer (pH 5.0) was dispensed into the wells and the plate was shaken

for two minutes (shaking diameter 1 mm, 1 020 rpm), after which luminescence was recorded with a measuring time of 1000 ms.

Plating

[0035] Samples for plating were diluted 10^2 to 10^7 fold with 150 mM NaCl and plated onto L agar plates (L broth containing 1.6 % agar). Colonies were counted after overnight incubation at 37 °C.

Live/dead staining and Flow cytometric analysis

[0036] Bacteria from 1 000 µl of cell culture were used for live/dead staining and flow cytometric analysis using a LIVE/DEAD BacLight bacterial viability kit (catalogue no. L-7005) for microscopy and quantitative analysis obtained from Molecular Probes Europe (Leiden, The Netherlands) and Fluoresbrite beads (diameter, 1.8 µm) obtained from Polysciences Inc. (Warrington, Pa.) as described in Virta et al. (1998) Appl. Environ. Microbiol. 64: 515–519.

Results

[0037] When the cultures were transferred to +30 °C, the cells grew logarithmically for 1-4 hours depending on the initial cell concentration. Luminescence and fluorescence rose logarithmically and were essentially constant per cell. Thus cell number could be assessed based on luminescence or fluorescence.

[0038] When ethanol was added in different concentrations to the growth medium (see Figures 4 and 5) death was, after a very short incubation period of 5 min, more or less insignificant at ethanol concentrations below 5 % and became more significant with increasing ethanol concentration reaching very pronounced significance at ethanol concentrations above 10 %. Correspondingly fluorescence (Figure 2) was essentially constant whatever the ethanol concentration in spite of dramatically decreasing corresponding live cell count according to plate count (Figure 4) and percentage of live cells according to the live/dead staining (Figure 5) whereas luminescence (Figure 3) dropped dramatically essentially corresponding to the dramatic drop in plate count (Figure 4) and the percentage of live cells (Figure 5) with increased ethanol concentration.

[0039] The effect of serum complement on the growth and death of *E. coli* is shown in Figures 6 to 8. Fluorescence (Figure 6) and luminescence (Figure 7) are shown before (squares) and after (circles) incubation for 90 minutes with serum complement. Fluorescence (Figure 6) is

slightly increased, during incubation regardless of the concentration of serum, whereas luminescence (Figure 7) decreases during incubation with increasing serum concentration. The decrease of luminescence during incubation with increasing concentrations of serum correlates clearly with the percentage of cells alive after incubation (Figure 8).

CLAIMS

1. A method to enable the assessment of the growth rate and death rate of a micro-organism within a chosen time period in an environment of interest by introducing into said micro-organism at least two reporter genes, which method is characterised in that
 - a) said reporter genes code for luminescent and/or fluorescent products and within said time period and environment at least two said products of the following are produced:
 - i) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that are or have been alive within said chosen time period,
 - ii) a product present in said environment of interest in an essentially known proportion to the amount of cells alive at any moment within said chosen time period, and
 - iii) an essentially stable product produced in a, within the environment of interest, essentially known proportion to the total amount of cells of said micro-organism that have died within said chosen time period,and said products can be measured through their luminescence and/or fluorescence;
 - b) said micro-organism is incubated within the environment of interest and said luminescence and/or fluorescence is detected after said chosen time period, and
 - c) the growth and death rate of the said micro-organism is assessed based on at least two of the following:
 - i) the known proportion of luminescence or fluorescence to the amount of cells alive after any said chosen time period,
 - ii) the known proportion of luminescence or fluorescence to the total amount of cells that are or have been alive within any said chosen time period, and
 - iii) the known proportion of luminescence or fluorescence to the total amount of cells that have died within any said chosen time period.
2. The method according to claim 1 characterised in that said micro-organism is a gram negative bacteria, e.g. *Escherichia coli*.

3. The method according to claim 1 characterised in that
- a) one reporter gene coding for a luminescent product is luciferase, which is used for the determination of amount of cells alive at any moment within said chosen time period, and
 - b) another reporter gene coding for a fluorescent product is green fluorescent protein (GFP), which is used for the determination of total amount of cells of said micro-organism that are or have been alive within said chosen time period.
4. The method according to claim 1 characterised in that said reporter genes are introduced into said micro-organism in a plasmid.
5. The method according to claim 3 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).
6. The method according to claim 2 characterised in that
- a) one reporter gene coding for a luminescent product is luciferase, which is used for the determination of amount of cells alive at any moment within said chosen time period, and
 - b) another reporter gene coding for a fluorescent product is green fluorescent protein (GFP), which is used for the determination of total amount of cells of said micro-organism that are or have been alive within said chosen time period.
7. The method according to claim 2 characterised in that said reporter genes are introduced into said micro-organism in a plasmid.
8. The method according to claim 4 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).
9. The method according to claim 6 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).

10. The method according to claim 7 characterised in that said plasmid is pGFP+luc* (SEQ ID NO: 1).

0980585-030402

ABSTRACT

[0040] A method to enable the assessment of the growth rate and death rate of a micro-organism within a chosen time period in an environment of interest. The method is characterised in that

5 a) two reporter genes are introduced to said micro-organism wherein, the reporter genes used code for luminescent and/or fluorescent products and at least two of the following products: an essentially stable product produced in an essentially known proportion to the total amount of cells of said micro-organism that are or have been alive within said chosen time period; a product present in an essentially known proportion to the amount of cells alive at any moment within said chosen time period; and an essentially stable product produced in an essentially known proportion to the total amount of cells of the said micro-organism that have died within said chosen time period, and said products can be measured through their luminescence and/or fluorescence;

10 b) the said micro-organism is incubated and said luminescence and/or fluorescence is detected after said chosen time periods, and

15 c) the growth and death rate of the said micro-organism is assessed based on at least two of the following: the known proportion of luminescence or fluorescence to the amount of cells alive after any said chosen time period; the known proportion of luminescence or fluorescence to the total amount of cells that are or have been alive within any said chosen time period; and the known proportion of luminescence or fluorescence to the total amount of cells that have died within any said chosen time period.




Figure 1

09980585-030402




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

201027-58508660




Figure 8

DECLARATION AND POWER OF ATTORNEY FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63)		Attorney Docket No.	
		First Named Inventor	
		COMPLETE IF KNOWN	
		Application Number	
		Filing Date	
		Group Art Unit	
<input type="checkbox"/> Declaration Submitted with Initial Filing	<input type="checkbox"/> Declaration Submitted after Initial Filing	Examiner Name	

As a below named inventor, I hereby declare that:

My residence, mailing address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: A method to enable assessment of growth and death of micro-organisms

☐ the specification of which is attached hereto.

☒ the specification of which was filed on December 4, 2001 as United States Application Number 09/980,585 or PCT International Application Number PCT/FI00/00507 and was amended on December 4, 2001

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Numbers	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
991296	FI	06/07/1999	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I or we hereby appoint the registered practitioner(s) associated with Customer No. **6449** to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Direct all correspondence to Customer Number **6449**.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name <u>Esa-Matti</u> (first and middle [if any])		Family Name <u>Lilius</u> or Surname	
Inventor's Signature <u>Esa-Matti</u>		Date <u>21.1.2002</u>	
Residence: City <u>Kaarina</u>	State <u>FIN</u>	Country <u>Finland</u>	Citizenship <u>Finnish</u>
Mailing Address <u>Vaakunatie 10</u>			
Mailing Address			
City <u>Kaarina</u>	State	Zip <u>FIN-20780</u>	Country <u>Finland</u>
NAME OF SECOND INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name <u>Marko</u> (first and middle [if any])		Family Name <u>Virta</u> or Surname	
Inventor's Signature <u>Marko</u>		Date <u>20.1.2002</u>	
Residence: City <u>Turku</u>	State	Country <u>Finland</u>	Citizenship <u>Finnish</u>
Mailing Address <u>Kauppiaskatu 10 D 59</u>			
Mailing Address			
City <u>Turku</u>	State	Zip <u>FIN-20100</u>	Country <u>Finland</u>
NAME OF THIRD INVENTOR:		<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name (first and middle [if any])		Family Name or Surname	
Inventor's Signature		Date	
Residence: City	State	Country	Citizenship
Mailing Address			
Mailing Address			
City	State	Zip	Country

SEQUENCE LISTING

<110> LILIIUS, Esa-Matti
VIRTA, Marko

<120> A Method to Enable Assessment of Growth and Death of Micro-organisms

<130> 2328-124

<150> PCT/FI00/00507
<151> 2000-06-07

<150> FI 991296
<151> 1999-06-07

<160> 4

<170> PatentIn version 3.0

<210> 1
<211> 5051
<212> DNA
<213> Artificial

<220>
<223> pGFP+Luc* plasmid

<220>
<221> promoter
<222> (95)..(199)
<223> lac promoter

<220>
<221> CDS
<222> (289)..(1005)
<223> GFP

<220>
<221> CDS
<222> (1044)..(2693)
<223> firefly luciferase

<220>
<221> CDS
<222> (3251)..(4108)
<223> beta-lactamase

<400> 1
agcgcccaat acgcaaacgc cctctccccc cgcggttgcc gattcattaa tgcagctggc 60
acgacagggt tcccgactgg aaagcgggca gtgagcgcaa cgcaattaat gtgagttagc 120
tcaactcatta ggcaccccag gctttacact ttatgcttcc ggctcgtagt ttgtgtggaa 180
ttgtgagcgg ataacaattt cacacaggaa acagctatga ccatgattac gccaaagcttg 240
catgcctgca ggtcgactct agaggatccc cgggtaccgg tcgccacc atg gtg agc 297
Met Val Ser
1
aag ggc gag gag ctg ttc acc ggg gtg gtg ccc atc ctg gtc gag ctg 345
Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu
5 10 15

0980585-1340

gac ggc gac gta aac ggc cac aag ttc agc gtg tcc ggc gag ggc gag 393
 Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu
 20 25 30 35

ggc gat gcc acc tac ggc aag ctg acc ctg aag ttc atc tgc acc acc 441
 Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr
 40 45 50

ggc aag ctg ccc gtg ccc tgg ccc acc ctc gtg acc acc ctg acc tac 489
 Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr
 55 60 65

ggc gtg cag tgc ttc agc cgc tac ccc gac cac atg aag cag cac gac 537
 Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp
 70 75 80

ttc ttc aag tcc gcc atg ccc gaa ggc tac gtc cag gag cgc acc atc 585
 Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Val Glu Arg Thr Ile
 85 90 95

ttc ttc aag gac gac ggc aac tac aag acc cgc gcc gag gtg aag ttc 633
 Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe
 100 105 110 115

gag ggc gac acc ctg gtg aac cgc atc gag ctg aag ggc atc gac ttc 681
 Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe
 120 125 130

aag gag gac ggc aac atc ctg ggc cac aag ctg gag tac aac tac aac 729
 Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn
 135 140 145

agc cac aac gtc tat atc atg gcc gac aag cag aag aac ggc atc aag 777
 Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys
 150 155 160

gtg aac ttc aag atc cgc cac aac atc gag gac ggc agc gtg cag ctc 825
 Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu
 165 170 175

gcc gac cac tac cag cag aac acc ccc atc ggc gac ggc ccc gtg ctg 873
 Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu
 180 185 190 195

ctg ccc gac aac cac tac ctg agc acc cag tcc gcc ctg agc aaa gac 921
 Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp
 200 205 210

ccc aac gag aag cgc gat cac atg gtc ctg ctg gag ttc gtg acc gcc 969
 Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala
 215 220 225

gcc ggc atc act ctc ggc atg gac gag ctg tac aag taaagcggcc 1015
 Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
 230 235

gctctagaac tagtgatcc ccctgacc atg gaa gac gcc aaa aac ata aag 1067
 Met Glu Asp Ala Lys Asn Ile Lys
 240 245

aaa ggc ccg gcg cca ttc tat ccg cta gag gat gga acc gct gga gag 1115
 Lys Gly Pro Ala Pro Phe Tyr Pro Leu Glu Asp Gly Thr Ala Gly Glu

000000.000000

250						255						260						
caa Gln	ctg Leu 265	cat His	aag Lys	gct Ala	atg Met	aag Lys 270	aga Arg	tac Tyr	gcc Ala	ctg Leu	gtt Val 275	cct Pro	gga Gly	aca Thr	att Ile	1163		
gct Ala 280	ttt Phe	aca Thr	gat Asp	gca Ala	cat His 285	atc Ile	gag Glu	gtg Val	aac Asn	atc Ile 290	acg Thr	tac Tyr	gcg Ala	gaa Glu	tac Tyr 295	1211		
ttc Phe	gaa Glu	atg Met	tcc Ser	gtt Val 300	cgg Arg	ttg Leu	gca Ala	gaa Glu	gct Ala 305	atg Met	aaa Lys	cga Arg	tat Tyr	ggg Gly 310	ctg Leu	1259		
aat Asn	aca Thr	aat Asn	cac His 315	aga Arg	atc Ile	gtc Val	gta Val	tcg Cys 320	agt Ser	gaa Glu	aac Asn	tct Ser	ctt Leu 325	caa Gln	ttc Phe	1307		
ttt Phe	atg Met	cgg Pro 330	gtg Val	ttg Leu	ggc Gly	cgc Ala	tta Leu 335	ttt Phe	atc Ile	gga Gly	gtt Val	gca Ala 340	gtt Val	gcg Ala	ccc Pro	1355		
gcg Ala 345	aac Asn	gac Asp	att Ile	tat Tyr	aat Asn	gaa Glu 350	cgt Arg	gaa Glu	ttg Leu	ctc Leu	aac Asn 355	agt Ser	atg Ser	aac Asn	att Ile	1403		
tcg Ser 360	cag Gln	cct Pro	acc Thr	gta Val	gtg Val 365	ttt Phe	gtt Val	tcc Ser	aaa Lys	aag Lys 370	ggg Gly	ttg Leu	caa Gln	aaa Lys	att Ile 375	1451		
ttg Leu	aac Asn	gtg Val	caa Gln	aaa Lys 380	aaa Lys	tta Leu	cca Pro	ata Ile	atc Ile 385	cag Gln	aaa Lys	att Ile	att Ile	atc Ile 390	atg Met	1499		
gat Asp	tct Ser	aaa Lys	acg Thr 395	gat Asp	tac Tyr	cag Gln	gga Gly	ttt Phe 400	cag Gln	tcg Ser	atg Met	tac Tyr	acg Thr 405	ttc Phe	gtc Val	1547		
aca Thr	tct Ser	cat His 410	cta Leu	cct Pro	ccc Pro	ggt Gly	ttt Phe 415	aat Asn	gaa Glu	tac Tyr	gat Asp	ttt Phe 420	gta Val	cca Pro	gag Glu	1595		
tcc Ser	ttt Phe 425	gat Asp	cgt Arg	gac Asp	aaa Lys	aca Thr 430	att Ile	gca Ala	ctg Leu	ata Ile	atg Met 435	aac Asn	tcc Ser	tct Ser	gga Gly	1643		
tct Ser 440	act Thr	ggg Gly	tta Leu	cct Pro	aag Lys 445	ggt Gly	gtg Val	gcc Ala	ctt Leu	ccg Pro 450	cat His	aga Arg	act Thr	gcc Ala	tgc Cys 455	1691		
gtc Val	aga Arg	tgc Phe	tcg Ser	cat His 460	gcc Ala	aga Arg	gat Asp	cct Pro	att Ile 465	ttt Phe	ggc Gly	aat Asn	caa Gln	atc Ile 470	att Ile	1739		
ccg Pro	gat Asp	act Thr 475	gcg Ala	att Ile	tta Leu	gta Ser	gtt Val	gta Val 480	cca Pro	ttc Phe	cat His	cac His	ggt Gly 485	ttt Phe	gga Gly	1787		
atg Met	ttt Phe	act Thr 490	aca Thr	ctc Gly	gga Gly	tat Tyr	ttg Leu 495	ata Ile	tgt Cys	gga Gly	ttt Phe	cga Arg 500	gtc Val	gtc Val	tta Leu	1835		
atg Met	tat Thr	aga Phe	gta Thr	gaa Glu	gaa Glu	gag Glu	ctg Leu	ttt Phe	tta Gly	cga Ser	tcc Leu	ctt Leu	cag Gln	gat Asp	tac Tyr	1883		

505					510					515						
aaa Lys 520	att Ile	caa Gln	agt Ser	gcy Ala	ttg Leu 525	cta Leu	gta Val	cca Pro	acc Thr	cta Leu 530	ttt Phe	tca Ser	ttc Phe	ttc Phe	gcc Ala 535	1931
aaa Lys	agc Ser	act Thr	ctg Leu	att Ile 540	gac Asp	aaa Lys	tac Tyr	gat Asp	tta Leu 545	tct Ser	aat Asn	tta Leu	cac His	gaa Glu 550	att Ile	1979
gct Ala	tct Ser	ggg Gly	ggc Gly 555	gca Ala	cct Pro	ctt Leu	tcg Ser	aaa Lys 560	gaa Glu	gtc Val	ggg Gly	gaa Glu	gcg Ala 565	ggt Val	gca Ala	2027
aaa Lys	cgc Arg	ttc Phe 570	cat His	ctt Leu	cca Pro	ggg Gly	ata Ile 575	cga Arg	caa Gln	gga Gly	tat Tyr	ggg Gly 580	ctc Leu	act Thr	gag Glu	2075
act Thr	aca Thr 585	tca Ser	gct Ala	att Ile	ctg Leu	att Ile 590	aca Thr	ccc Pro	gag Glu	ggg Gly	gat Asp 595	gat Asp	aaa Lys	ccg Pro	ggc Gly	2123
gcg Ala 600	gtc Val	ggt Gly	aaa Lys	gtt Val 605	cca Pro	ttt Phe	ttt Phe	gaa Glu	gcg Ala 610	aag Lys	gtt Val	gtg Val	gat Asp	ctg Leu 615		2171
gat Asp	acc Thr	ggg Gly	aaa Lys	acg Thr 620	ctg Leu	ggc Gly	gtt Val	aat Asn	cag Gln 625	aga Arg	ggc Gly	gaa Glu	tta Leu	tgt Cys 630	gtc Val	2219
aga Arg	gga Gly	cct Pro	atg Met 635	att Ile	atg Met	tcc Ser	ggt Gly	tat Tyr 640	gta Val	aac Asn	aat Asn	ccg Pro	gaa Glu 645	gcg Ala	acc Thr	2267
aac Asn	gcc Ala	ttg Leu 650	att Ile	gac Asp	aag Lys	gat Asp 655	gga Gly	tgg Trp	cta Leu	cat His	tct Ser	gga Gly 660	gac Asp	ata Ile	gct Ala	2315
tac Tyr	tgg Trp 665	gac Asp	gaa Glu	gac Asp	gaa Glu	cac His 670	ttc Phe	ttc Phe	ata Ile	gtt Val	gac Asp 675	cgc Arg	ttg Leu	aag Lys	tct Ser	2363
tta Leu 680	att Ile	aaa Lys	tac Tyr	aaa Lys	gga Gly 685	tac Tyr	cag Gln	gtg Val	gcc Ala 690	ccc Pro	gct Ala	gaa Glu	ttg Leu	gag Glu	tcg Ser 695	2411
ata Ile	ttg Leu	tta Leu	caa Gln	cac His 700	ccc Pro	aac Asn	atc Ile	ttc Phe	gac Asp 705	gcg Ala	ggc Gly	gtg Val	gca Ala 710	ggt Gly	ctt Leu	2459
ccc Pro	gac Asp	gat Asp	gac Asp 715	gcc Ala	ggg Gly	gaa Glu	ctt Leu	ccc Pro 720	gcc Ala	gcc Ala	ggt Val	ggt Val	ggt Val 725	ttg Leu	gag Glu	2507
cac His	gga Gly	aag Lys 730	acg Thr	atg Met	acg Thr	gaa Glu	aaa Lys 735	gag Glu	atc Ile	gtg Val	gat Asp	tac Tyr 740	gtc Val	gcc Ala	agt Ser	2555
caa Gln	gta Val 745	aca Thr	acc Thr	gcc Ala	aaa Lys	aag Lys 750	ttg Leu	cgc Arg	gga Gly	gga Gly	ggt Val 755	gtg Val	ttt Phe	gtg Val	gac Asp	2603
gaa Gln	gta Val	ccg Arg	aaa Lys	ggg Gly	ctt Leu	acc Pro	gga Glu	aaa Lys	ctc Gln	gac Ala	gca Ala	aga Gln	aaa Lys	atc Gln	aga Gln	2651

Glu Val Pro Lys Gly Leu Thr Gly Lys Leu Asp Ala Arg Lys Ile Arg	760	765	770	775	
gag atc ctc ata aag gcc aag aag ggc gga aag tcc aaa ttg					2693
Glu Ile Leu Ile Lys Ala Lys Lys Gly Gly Lys Ser Lys Leu	780		785		
taaaaatgtaa ctgtattcag cgatgacgaa attcttagctt attgtaatac tctaggggct					2753
gcaggaattc gatatcaagc ttatcgatac cgtcgacctc gagggggggc cctttcgtct					2813
cgcgcggttc ggtgatgacg gtgaaaacct ctgacacatg cagctcccgct agacgggtcac					2873
agctttgtctg taagcggatg ccggggagcag acaagcccgt caggggcgct cagcgggtgt					2933
tggcgggtgt cggggctggc ttaactatgc ggcatcagag cagattgtac tgagagtgca					2993
ccatatgcgg tgtgaaatac cgcacagatg cgttaaggaga aaataccgca tcaggcggcc					3053
ttaagggcct cgtgatacgc ctatttttat aggttaatgt catgataata atggttttctt					3113
agaagtcagg tggcactttt cggggaaatg tgcgcgggaac ccctatttgt ttatttttct					3173
aaatacattc aaatatgtat ccgctcatga gacaataacc ctgataaatg cttcaataat					3233
attgaaaaag gaagagt atg agt att caa cat ttc cgt gtc gcc ctt att					3283
Met Ser Ile Gln His Phe Arg Val Ala Leu Ile	790		795	800	
ccc ttt ttt gcg gca ttt tgc ctt cct gtt ttt gct cac cca gaa acg					3331
Pro Phe Phe Ala Ala Phe Cys Leu Pro Val Phe Ala His Pro Glu Thr	805		810	815	
ctg gtg aaa gta aaa gat gct gaa gat cag ttg ggt gca cga gtg ggt					3379
Leu Val Lys Val Lys Asp Ala Glu Asp Gln Leu Gly Ala Arg Val Gly	820		825	830	
tac atc gaa ctg gat ctc aac agc ggt aag atc ctt gag agt ttt cgc					3427
Tyr Ile Glu Leu Asp Leu Asn Ser Gly Lys Ile Leu Leu Ser Phe Arg	835		840	845	
ccc gaa gaa cgt ttt cca atg atg agc act ttt aaa gtt ctg cta tgt					3475
Pro Glu Glu Arg Phe Pro Met Met Ser Thr Phe Lys Val Leu Leu Cys	850		855	860	
ggc gcg gta tta tcc cgt att gac gcc ggg caa gag caa ctc ggt cgc					3523
Gly Ala Val Leu Ser Arg Ile Asp Ala Gly Lys Glu Gln Leu Gly Arg	865		870	875	880
cgc ata cac tat tct cag aat gac ttg gtt gag tac tca cca gtc aca					3571
Arg Ile His Tyr Ser Gln Asn Asp Leu Val Glu Tyr Ser Pro Val Thr	885		890	895	
gaa aag cat ctt acg gat ggc atg aca gta aga gaa tta tgc agt gct					3619
Glu Lys His Leu Thr Asp Gly Met Thr Val Arg Glu Leu Cys Ser Ala	900		905	910	
gcc ata acc atg agt gat aac act gcg gcc aac tta ctt ctg aca acg					3667
Ala Ile Thr 915 Met Ser Asp Asn Thr Ala Ala Asn Leu Leu Thr Thr			920	925	
atc gga gga cgc aag gag cta acc gct ttt ttg cac aca atg ggg gat					3715
Ile Gly Gly Pro Lys Glu Leu Thr Ala Phe Leu His Asn Met Gly Asp					

4978

5038

5051

<211> 239

<212> PRT

<213> Artificial

<400> 2

Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
1 5 10 15

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
20 25 30

Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
35 40 45

Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
50 55 60

Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
65 70 75 80

Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85 90 95

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100 105 110

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115 120 125

```

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
  130                               135             140

```

Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145 150 155 160

Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165 170 175

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180 185 190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195 200 205

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210 215 220

Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
225 230 235

 $\langle 210 \rangle$ 3 $\langle 211 \rangle$ 550

<212> PRT

<213> Artificial

<400> 3

Met	Glu	Asp	Ala	Lys	Asn	Ile	Lys	Lys	Gly	Pro	Ala	Pro	Phe	Tyr	Pro
1			5						10					15	
Leu	Glu	Asp	Gly	Thr	Ala	Gly	Glu	Gln	Leu	His	Lys	Ala	Met	Lys	Arg
			20					25					30		
Tyr	Ala	Leu	Val	Pro	Gly	Thr	Ile	Ala	Phe	Thr	Asp	Ala	His	Ile	Glu
		35					40					45			
Val	Asn	Ile	Thr	Tyr	Ala	Glu	Tyr	Phe	Glu	Met	Ser	Val	Arg	Leu	Ala
		50				55					60				
Glu	Ala	Met	Lys	Arg	Tyr	Gly	Leu	Asn	Thr	Asn	His	Arg	Ile	Val	Val
65					70					75				80	
Cys	Ser	Glu	Asn	Ser	Leu	Gln	Phe	Phe	Met	Pro	Val	Leu	Gly	Ala	Leu
			85						90					95	
Phe	Ile	Gly	Val	Ala	Val	Ala	Pro	Ala	Asn	Asp	Ile	Tyr	Asn	Glu	Arg
			100					105					110		
Glu	Leu	Leu	Asn	Ser	Met	Asn	Ile	Ser	Gln	Pro	Thr	Val	Val	Phe	Val
		115					120					125			
Ser	Lys	Lys	Gly	Leu	Gln	Lys	Ile	Leu	Asn	Val	Gln	Lys	Lys	Leu	Pro
		130				135						140			
Ile	Ile	Gln	Lys	Ile	Ile	Ile	Met	Asp	Ser	Lys	Thr	Asp	Tyr	Gln	Gly
145					150					155				160	
Phe	Gln	Ser	Met	Tyr	Thr	Phe	Val	Thr	Ser	His	Leu	Pro	Pro	Gly	Phe
				165					170					175	
Asn	Glu	Tyr	Asp	Phe	Val	Pro	Glu	Ser	Phe	Asp	Arg	Asp	Lys	Thr	Ile
			180					185					190		
Ala	Leu	Ile	Met	Asn	Ser	Ser	Gly	Ser	Thr	Gly	Leu	Pro	Lys	Gly	Val
			195				200					205			
Ala	Leu	Pro	His	Arg	Thr	Ala	Cys	Val	Arg	Phe	Ser	His	Ala	Arg	Asp
					215						220				
Pro	Ile	Phe	Gly	Asn	Gln	Ile	Ile	Pro	Asp	Thr	Ala	Ile	Leu	Ser	Val
225					230					235					240
Val	Pro	Phe	His	His	Gly	Phe	Gly	Met	Phe	Thr	Thr	Leu	Gly	Tyr	Leu
				245					250					255	
Ile	Cys	Gly	Phe	Phe	Arg	Val	Val	Leu	Met	Tyr	Arg	Phe	Glu	Glu	Leu
			260						265				270		
Phe	Leu	Arg	Ser	Leu	Gln	Asp	Tyr	Lys	Ile	Gln	Ser	Ala	Leu	Leu	Val
		275					280					285			
Pro	Thr	Leu	Phe	Ser	Phe	Phe	Ala	Lys	Ser	Thr	Leu	Ile	Asp	Lys	Tyr
		290				295					300				
Asp	Leu	Ser	Asn	Leu	His	Glu	Ile	Ala	Ser	Gly	Gly	Ala	Pro	Leu	Ser
305					310					315					320
Lys	Glu	Val	Gly	Glu	Ala	Val	Ala	Lys	Arg	Phe	His	Leu	Pro	Gly	Ile
				325					330					335	
Arg	Gln	Gly	Tyr	Gly	Leu	Thr	Glu	Thr	Thr	Ser	Ala	Ile	Leu	Ile	Thr

340 345 350
 Pro Glu Gly Asp Asp Lys Pro Gly Ala Val Gly Lys Val Val Pro Phe
 355 360 365
 Phe Glu Ala Lys Val Val Asp Leu Asp Thr Gly Lys Thr Leu Gly Val
 370 375 380
 Asn Gln Arg Gly Glu Leu Cys Val Arg Gly Pro Met Ile Met Ser Gly
 385 390 395 400
 Tyr Val Asn Asn Pro Glu Ala Thr Asn Ala Leu Ile Asp Lys Asp Gly
 405 410 415
 Trp Leu His Ser Gly Asp Ile Ala Tyr Trp Asp Glu Asp Glu His Phe
 420 425 430
 Phe Ile Val Asp Arg Leu Lys Ser Leu Ile Lys Tyr Lys Gly Tyr Gln
 435 440 445
 Val Ala Pro Ala Glu Leu Glu Ser Ile Leu Leu Gln His Pro Asn Ile
 450 455 460
 Phe Asp Ala Gly Val Ala Gly Leu Pro Asp Asp Ala Gly Glu Leu
 465 470 475 480
 Pro Ala Ala Val Val Val Leu Glu His Gly Lys Thr Met Thr Glu Lys
 485 490 495
 Glu Ile Val Asp Tyr Val Ala Ser Gln Val Thr Thr Ala Lys Lys Leu
 500 505 510
 Arg Gly Gly Val Val Phe Val Asp Glu Val Pro Lys Gly Leu Thr Gly
 515 520 525
 Lys Leu Asp Ala Arg Lys Ile Arg Glu Ile Leu Ile Lys Ala Lys Lys
 530 535 540
 Gly Gly Lys Ser Lys Leu
 545 550

 <210> 4
 <211> 286
 <212> PRT
 <213> Artificial
 <400> 4
 Met Ser Ile Gln His Phe Arg Val Ala Leu Ile Pro Phe Phe Ala Ala
 1 5 10 15
 Phe Cys Leu Pro Val Phe Ala His Pro Glu Thr Leu Val Lys Val Lys
 20 25 30
 Asp Ala Glu Asp Gln Leu Gly Ala Arg Val Gly Tyr Ile Glu Leu Asp
 35 40 45
 Leu Asn Ser Gly Lys Ile Leu Glu Ser Phe Arg Pro Glu Glu Arg Phe
 50 55 60
 Pro Met Met Ser Thr Phe Lys Val Leu Leu Cys Gly Ala Val Leu Ser
 65 70 75 80
 Arg Ile Asp Ala Gly Gln Glu Gln Leu Gly Arg Arg Ile His Tyr Ser
 85 90 95

20250515.030402

Gln	Asn	Asp	Leu	Val	Glu	Tyr	Ser	Pro	Val	Thr	Glu	Lys	His	Leu	Thr
			100					105					110		
Asp	Gly	Met	Thr	Val	Arg	Glu	Leu	Cys	Ser	Ala	Ala	Ile	Thr	Met	Ser
		115					120					125			
Asp	Asn	Thr	Ala	Ala	Asn	Leu	Leu	Leu	Thr	Thr	Ile	Gly	Gly	Pro	Lys
		130				135					140				
Glu	Leu	Thr	Ala	Phe	Leu	His	Asn	Met	Gly	Asp	His	Val	Thr	Arg	Leu
145					150					155					160
Asp	Arg	Trp	Glu	Pro	Glu	Leu	Asn	Glu	Ala	Ile	Pro	Asn	Asp	Glu	Arg
				165					170					175	
Asp	Thr	Thr	Met	Pro	Val	Ala	Met	Ala	Thr	Thr	Leu	Arg	Lys	Leu	Leu
			180					185					190		
Thr	Gly	Glu	Leu	Leu	Thr	Leu	Ala	Ser	Arg	Gln	Gln	Leu	Ile	Asp	Trp
		195					200					205			
Met	Glu	Ala	Asp	Lys	Val	Ala	Gly	Pro	Leu	Leu	Arg	Ser	Ala	Leu	Pro
		210				215					220				
Ala	Gly	Trp	Phe	Ile	Ala	Asp	Lys	Ser	Gly	Ala	Gly	Glu	Arg	Gly	Ser
225					230					235					240
Arg	Gly	Ile	Ile	Ala	Ala	Leu	Gly	Pro	Asp	Gly	Lys	Pro	Ser	Arg	Ile
			245						250					255	
Val	Val	Ile	Tyr	Thr	Thr	Gly	Ser	Gln	Ala	Thr	Met	Asp	Glu	Arg	Asn
			260					265					270		
Arg	Gln	Ile	Ala	Glu	Ile	Gly	Ala	Ser	Leu	Ile	Lys	His	Trp		
		275					280					285			

SEQUENCE LISTING

<110> LILIUS, Esa-Matti
VIRTA, Marko

<120> A Method to Enable Assessment of Growth and Death of Micro-organisms

<130> 2328-124

<150> PCT/FI00/00507
<151> 2000-06-07

<150> FI 991296
<151> 1999-06-07

<160> 4

<170> PatentIn version 3.0

<210> 1
<211> 5051
<212> DNA
<213> Artificial

<220>
<223> pGFP+Luc* plasmid

<220>
<221> promoter
<222> (95)..(199)
<223> lac promoter

<220>
<221> gene
<222> (289)..(1005)
<223> Coding sequence for GFP

<220>
<221> gene
<222> (1044)..(2693)
<223> coding sequence for firefly luciferase

<220>
<221> gene
<222> (3251)..(4108)
<223> coding sequence for beta-lactamase

<400> 1
agcgcccaat acgcaaacgg cctctccccg cgcgttggcc gattcattaa tgcagctggc 60
acgacagggt tcccgactgg aaagcgggca gtgagcgcaa cgcaattaat gtgagttagc 120
tcactcatta ggcaccccgag gctttacact ttatgcttcc ggctcgatg ttgtgtggaa 180
ttgtgagcgg ataacaattt cacacaggaa acagctatga ccatgattac gccaagett 240
catgcttgca ggctgactct agaggatccc cgggtaccgg tcgccacat ggtgagcaag 300
ggcgaggagc tgttcaccgg ggtggtgccc atcctggtcg agctggacgg cgacgtaaac 360

ggccacaagt	tcagcgtgtc	cggcgagggc	gagggcgatg	ccacctacgg	caagctgacc	420
ctgaagttca	tctgccacc	cggcaagctg	cccgctgcct	ggcccacct	cgtgaccacc	480
ctgacctacg	gcgtgcagtg	cttcagcgc	taccocgacc	acatgaagca	gcacgacttc	540
ttcaagttcg	ccatgcccga	aggctactgc	caggagcgca	ccatcttctt	caaggaagac	600
ggcaactaca	agaccgcgc	cgaggtgaag	ttcgaggcgc	acacctggt	gaaccgcac	660
gagctgaagg	gcctgcactt	caaggaggac	ggcaacatcc	tggggcaca	gctggagtac	720
aactacaaca	gccacaacgt	ctatatcatg	gccgacaagc	agaagaacgg	catcaaggtg	780
aacttcaaga	tcgccacaa	catcgaggac	ggcagcgtgc	agctcgccga	ccactaccag	840
cagaacaccc	ccatcgccga	cggccccgtg	ctgctgccg	acaaccacta	cctgagcacc	900
cagtccgcc	tgagcaaaag	ccccaacgag	aagcgcgac	acatggtct	gctggagttc	960
gtgaccgcgc	ccgggatcac	tctcgcgatg	gacgagctgt	acaagtaaa	cggccgctct	1020
agaactagtg	gatccccgt	accatggaag	acgccccaaa	cataaagaaa	ggcccgccgc	1080
cattctatcc	gctagaggat	ggaaccgcgt	gagagcaact	gcataaggct	atgaagagat	1140
acgccctggt	tctggaaca	attgctttta	cagatgcaca	tatcgaggtg	aacatcacgt	1200
acgcggaata	cttcgaaatg	tccgttcggt	tggcagaagc	tatgaaacga	tatgggctga	1260
atacaaatca	cagaatcgtc	gtatgcagtg	aaaactctct	tcaattcttt	atgccggtgt	1320
tggcgcgctt	atttatcgga	gttgcatgtg	cgcgcggaaa	cgacatttat	aatgaacgtg	1380
aattgctcaa	cagtatgaac	atttcgcagc	ctaccgtagt	gtttgtttcc	aaaaagggggt	1440
tgcaaaaaat	ttgaaacgtg	caaaaaaat	taccaataat	ccagaaaatt	attatcatgg	1500
attctaaaa	ggattaccag	ggatttcagt	cgatgtacac	gttcgtcaca	tctcatctac	1560
ctcccggttt	taatgaatac	gattttgtac	cagagtcctt	tgatcgtgac	aaaacaattg	1620
cactgataat	gaactcctct	ggatctactg	ggttacctaa	gggtgtggcc	cttcgcgata	1680
gaactgcctg	cgtcagattc	tcgcatgcc	gagatcctat	ttttggcaat	caaatcattc	1740
cggactactgc	gattttaaagt	gttggtccat	tccatcacgg	ttttggaatg	ttactacac	1800
tcgggatattt	gatattgtgga	tttcgagctg	tcttaagtga	tagatttgaa	gaagagctgt	1860
ttttacgac	ccttcaggat	tacaaaattc	aaagtgcgtt	gctagtacca	acctattttt	1920
cattctctgc	caaaagcact	ctgattgaca	aatacgattt	atctaattta	cacgaaaattg	1980
cttctggggg	cgcacctctt	tcgaaaagag	tcggggaagc	ggttgcaaaa	cgttccatc	2040
ttccagggat	acgacaagga	tatgggtcca	ctgagactac	atcagctatt	ctgattacac	2100
cggaggggga	tgataaacgg	ggcgcggctg	gtaaaagttg	tccatttttt	gaagcgaaag	2160

ttgtggatct	ggataccggg	aaaaacgtgg	gcgttaatca	gagaggogaa	ttatgtgtca	2220
gaggacctat	gattatgtcc	ggttatgtaa	acaatccgga	agcgaccaac	gccttgattg	2280
acaaggatgg	atggctacat	tctggagaca	tagcttactg	ggacgaagac	gaacaattct	2340
tcatagttda	ccgcttgaag	tctttaatta	aatacaaaag	ataccagggtg	gcccccgctg	2400
aattggagtc	gatattgtta	caacacccca	acatcttcga	cgcgggcgtg	gcagggtcttc	2460
ccgacgatga	cgccgggtga	cttcccgccg	ccgttgttgt	tttgaggcac	ggaaagacga	2520
tgacggaaaa	agagatcgtg	gattacgtcg	ccagtcaagt	aacaaccgcc	aaaaagttgc	2580
gcggaggagt	tgtgtttgtg	gacgaagtac	cgaaaggctc	taccggaaaa	ctcgacgcaa	2640
gaaaaatcag	agagatcctc	ataaaggcca	agaaggcgcg	aaagtccaaa	ttgtaaaaatg	2700
taactgtatt	cagcgatgac	gaaattctta	gctattgtaa	tactctaggg	gctgcaggaa	2760
ttcgatatca	agcttatcga	taccgtcgac	ctcgaggggg	ggccctttcg	tctcgccgct	2820
ttcggtgatg	acggtgaaaa	cctctgacac	atgcagctcc	cggagacggt	cacagcttgt	2880
ctgtaagcgg	atgccgggag	cagacaagcc	cgtcaggggc	cgtcagcggg	tggtggcggg	2940
tgtcggggct	ggcttaacta	tgccgcatca	gagcagattg	tactgagagt	gcaccatatg	3000
cgggtgtgaa	taccgcacag	atgcgtaaag	agaaaatacc	gcatacggcg	gccttaaggg	3060
cctcgtgata	cgctatcttt	tatagggtaa	tgctcatgata	ataatggttt	cttagacgtc	3120
agggtggcact	tttcggggaa	atgtgcgcgg	aaccctcttt	tggttttttt	tctaaataca	3180
ttcaaatatg	tatccgctca	tgagacaata	accctgataa	atgcttcaat	aatattgaaa	3240
aaggaaagat	atgagtattc	aacatttccg	tgctgcctct	attccctttt	ttgcggcatt	3300
ttgccttctc	gtttttgtgc	accagaaaaa	gctggtgaaa	gtaaaagatg	ctgaagatca	3360
gttgggtgca	cgagtgggtt	acatcgaaat	ggatctcaac	agcggtaaga	tccttgagag	3420
ttttgcctcc	gaagaacggt	ttccaatgat	gagcactttt	aaagtctcgc	tatgtggcgc	3480
ggtattatcc	cgtattgacg	ccgggcaaga	gcaactcggg	cgccgcatac	actattctca	3540
gaatgacttg	gttgagtact	caccagtcac	agaaaagcat	cttacgggatg	gcatagacagt	3600
aagagaatta	tgacgtgctg	ccataaccat	gagtgataac	actgcggcca	acttacttct	3660
gacacagatc	ggaggaccga	aggagctaac	cgcttttttg	cacaacatgg	gggatcatgt	3720
aaactgcctt	gatcgttggt	aacggagctg	gaatgaagcc	ataccaaacg	acgagcgtga	3780
caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaaa	ctattaactg	gcgaactact	3840
tactctagct	tcocggcaac	aattaataga	ctggatggag	gcggataaag	ttgcaggacc	3900
actctcgcgc	tcggcccttc	cggctggctg	gtttattgct	gataaatctg	gagccgggtga	3960

gcgtgggtct cgcggtatca ttgcagcact gggggccagat ggtaagccct cccgtatcgt 4020
 agttatctac acgacgggga gtcaggcaac tatggatgaa cgaaatagac agatcgctga 4080
 gataggtgcc tcaactgatta agcattggtta actgtcagac caagtttact catatatact 4140
 ttagattgat ttaaaacttc atttttaatt taaaaggatc taggtgaaga tcctttttga 4200
 taatctcatg accaaaaatcc cttaacgtga gttttcgttc cactgagcgt cagaccccggt 4260
 agaaaagatc aaaggatctt ctlgagatcc tttttttctg cgcgtaatct gctgcttgca 4320
 aacaaaaaaa ccaccgctac cagcgggtgtt ttgtttgccg gatcaagagc taccaactct 4380
 ttttccgaag gtaactgggt tcagcagagc gcagatacca aatactgtcc ttctagtgtta 4440
 gccgtagtta gcccaccact tcaagaactc tgtagcaccg cctacatacc tcgctctgct 4500
 aatcctgtta ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg ggttggaactc 4560
 aagacgatag ttaccggata aggcgcagcg gtcgggtgta acgggggggt cgtgcacaca 4620
 gccacgttg gagcgaacga cctacaccga actgagatac ctacagcgtg agctatgaga 4680
 aagcgccacg ctctccgaag ggagaaagcg ggacaggatc ccggtaaagc gcagggtcgg 4740
 aacaggagag cgcacgaggg agcttccagg gggaaacgcc tggtatcttt atagtctgt 4800
 cgggtttcgc caoctctgac ttgagcgtcg atttttgtga tgctcgtcag gggggcgagg 4860
 cctatggaaa aacgccagca acgcgcctt ttacggttc ctggcctttt gctggccttt 4920
 tgctcactg ttctttctcg cgttatcccc tgattctgtg gataaccgta ttaccgcctt 4980
 tgagtgaagt gataccgctc gccgcagccg aacgaccgag cgcagcgagt cagtgaagca 5040
 ggaagcgga g 5051

<210> 2
 <211> 239
 <212> PRT
 <213> Unknown

<220>
 <223> source of GFP unknown

<400> 2
 Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 1 5 10 15
 Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
 20 25 30
 Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
 35 40 45
 Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
 50 55 60

Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
65 70 75 80

Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85 90 95

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100 105 110

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115 120 125

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130 135 140

Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145 150 155 160

Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165 170 175

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180 185 190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195 200 205

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210 215 220

Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
225 230 235

<210> 3

<211> 550

<212> PRT

<213> Unknown

<220>

<223> source of firefly luciferase unknown

<400> 3

Met Glu Asp Ala Lys Asn Ile Lys Lys Gly Pro Ala Pro Phe Tyr Pro
1 5 10 15

Leu Glu Asp Gly Thr Ala Gly Glu Gln Leu His Lys Ala Met Lys Arg
20 25 30

Tyr Ala Leu Val Pro Gly Thr Ile Ala Phe Thr Asp Ala His Ile Glu
35 40 45

Val Asn Ile Thr Tyr Ala Glu Tyr Phe Glu Met Ser Val Arg Leu Ala
50 55 60

Glu Ala Met Lys Arg Tyr Gly Leu Asn Thr Asn His Arg Ile Val Val
65 70 75 80

Cys Ser Glu Asn Ser Leu Gln Phe Phe Met Pro Val Leu Gly Ala Leu
85 90 95

Phe Ile Gly Val Ala Val Ala Pro Ala Asn Asp Ile Tyr Asn Glu Arg
 100 105 110
 Glu Leu Leu Asn Ser Met Asn Ile Ser Gln Pro Thr Val Val Phe Val
 115 120 125
 Ser Lys Lys Gly Leu Gln Lys Ile Leu Asn Val Gln Lys Lys Leu Pro
 130 135 140
 Ile Ile Gln Lys Ile Ile Ile Met Asp Ser Lys Thr Asp Tyr Gln Gly
 145 150 155 160
 Phe Gln Ser Met Tyr Thr Phe Val Thr Ser His Leu Pro Pro Gly Phe
 165 170 175
 Asn Glu Tyr Asp Phe Val Pro Glu Ser Phe Asp Arg Asp Lys Thr Ile
 180 185 190
 Ala Leu Ile Met Asn Ser Ser Gly Ser Thr Gly Leu Pro Lys Gly Val
 195 200 205
 Ala Leu Pro His Arg Thr Ala Cys Val Arg Phe Ser His Ala Arg Asp
 210 215 220
 Pro Ile Phe Gly Asn Gln Ile Ile Pro Asp Thr Ala Ile Leu Ser Val
 225 230 235 240
 Val Pro Phe His His Gly Phe Gly Met Phe Thr Thr Leu Gly Tyr Leu
 245 250 255
 Ile Cys Gly Phe Arg Val Val Leu Met Tyr Arg Phe Glu Glu Glu Leu
 260 265 270
 Phe Leu Arg Ser Leu Gln Asp Tyr Lys Ile Gln Ser Ala Leu Leu Val
 275 280 285
 Pro Thr Leu Phe Ser Phe Phe Ala Lys Ser Thr Leu Ile Asp Lys Tyr
 290 295 300
 Asp Leu Ser Asn Leu His Glu Ile Ala Ser Gly Gly Ala Pro Leu Ser
 305 310 315 320
 Lys Glu Val Gly Glu Ala Val Ala Lys Arg Phe His Leu Pro Gly Ile
 325 330 335
 Arg Gln Gly Tyr Gly Leu Thr Glu Thr Thr Ser Ala Ile Leu Ile Thr
 340 345 350
 Pro Glu Gly Asp Asp Lys Pro Gly Ala Val Gly Lys Val Val Pro Phe
 355 360 365
 Phe Glu Ala Lys Val Val Asp Leu Asp Thr Gly Lys Thr Leu Gly Val
 370 375 380
 Asn Gln Arg Gly Glu Leu Cys Val Arg Gly Pro Met Ile Met Ser Gly
 385 390 395 400
 Tyr Val Asn Asn Pro Glu Ala Thr Asn Ala Leu Ile Asp Lys Asp Gly
 405 410 415

Trp Leu His Ser Gly Asp Ile Ala Tyr Trp Asp Glu Asp Glu His Phe
420 425 430

Phe Ile Val Asp Arg Leu Lys Ser Leu Ile Lys Tyr Lys Gly Tyr Gln
435 440 445

Val Ala Pro Ala Glu Leu Glu Ser Ile Leu Leu Gln His Pro Asn Ile
450 455 460

Phe Asp Ala Gly Val Ala Gly Leu Pro Asp Asp Ala Gly Glu Leu
465 470 475 480

Pro Ala Ala Val Val Val Leu Glu His Gly Lys Thr Met Thr Glu Lys
485 490 495

Glu Ile Val Asp Tyr Val Ala Ser Gln Val Thr Thr Ala Lys Lys Leu
500 505 510

Arg Gly Gly Val Val Phe Val Asp Glu Val Pro Lys Gly Leu Thr Gly
515 520 525

Lys Leu Asp Ala Arg Lys Ile Arg Glu Ile Leu Ile Lys Ala Lys Lys
530 535 540

Gly Gly Lys Ser Lys Leu
545 550

<210> 4
<211> 286
<212> PRT
<213> Unknown

<220>
<223> source of beta-lactamase unknown

<400> 4
Met Ser Ile Gln His Phe Arg Val Ala Leu Ile Pro Phe Phe Ala Ala
1 5 10 15

Phe Cys Leu Pro Val Phe Ala His Pro Glu Thr Leu Val Lys Val Lys
20 25 30

Asp Ala Glu Asp Gln Leu Gly Ala Arg Val Gly Tyr Ile Glu Leu Asp
35 40 45

Leu Asn Ser Gly Lys Ile Leu Glu Ser Phe Arg Pro Glu Glu Arg Phe
50 55 60

Pro Met Met Ser Thr Phe Lys Val Leu Leu Cys Gly Ala Val Leu Ser
65 70 75 80

Arg Ile Asp Ala Gly Gln Glu Gln Leu Gly Arg Arg Ile His Tyr Ser
85 90 95

Gln Asn Asp Leu Val Glu Tyr Ser Pro Val Thr Glu Lys His Leu Thr
100 105 110

Asp Gly Met Thr Val Arg Glu Leu Cys Ser Ala Ala Ile Thr Met Ser
115 120 125

Asp Asn Thr Ala Ala Asn Leu Leu Thr Thr Ile Gly Gly Pro Lys
130 135 140

Glu Leu Thr Ala Phe Leu His Asn Met Gly Asp His Val Thr Arg Leu
145 150 155 160

Asp Arg Trp Glu Pro Glu Leu Asn Glu Ala Ile Pro Asn Asp Glu Arg
165 170 175

Asp Thr Thr Met Pro Val Ala Met Ala Thr Thr Leu Arg Lys Leu Leu
180 185 190

Thr Gly Glu Leu Leu Thr Leu Ala Ser Arg Gln Gln Leu Ile Asp Trp
195 200 205

Met Glu Ala Asp Lys Val Ala Gly Pro Leu Leu Arg Ser Ala Leu Pro
210 215 220


Ala Gly Trp Phe Ile Ala Asp Lys Ser Gly Ala Gly Glu Arg Gly Ser
225 230 235 240

Arg Gly Ile Ile Ala Ala Leu Gly Pro Asp Gly Lys Pro Ser Arg Ile
245 250 255

Val Val Ile Tyr Thr Thr Gly Ser Gln Ala Thr Met Asp Glu Arg Asn
260 265 270

Arg Gln Ile Ala Glu Ile Gly Ala Ser Leu Ile Lys His Trp
275 280 285

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



SCANNED, # 12

Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ **Scanned copy is best available.**

- Color case only, 5 Drawing not 8.